



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

LIM Homeodomain Transcription Factor Isl-1 Enhances Follicle Stimulating Hormone-beta and Luteinizing Hormone-beta Gene Expression and Mediates the Activation of Leptin on Gonadotropin Synthesis

Citation for published version:

Wu, Y, Luo, H, Liu, J, Kang, D, McNeilly, AS & Cui, S 2010, 'LIM Homeodomain Transcription Factor Isl-1 Enhances Follicle Stimulating Hormone-beta and Luteinizing Hormone-beta Gene Expression and Mediates the Activation of Leptin on Gonadotropin Synthesis', *Endocrinology*, vol. 151, no. 10, pp. 4787-4800.
<https://doi.org/10.1210/en.2010-0208>

Digital Object Identifier (DOI):

[10.1210/en.2010-0208](https://doi.org/10.1210/en.2010-0208)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Endocrinology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



LIM Homeodomain Transcription Factor Isl-1 Enhances Follicle Stimulating Hormone- β and Luteinizing Hormone- β Gene Expression and Mediates the Activation of Leptin on Gonadotropin Synthesis

Yingjie Wu, Haoshu Luo, Jiali Liu, Duo Kang, Alan S. McNeilly, and Sheng Cui

State Key Laboratory of Agrobiotechnology (Y.W., H.L., J.L., D.K., S.C.), College of Biological Sciences, China Agricultural University, Beijing 100193, People's Republic of China; and Medical Research Council Human Reproductive Sciences Unit (A.S.M.), Queen's Medical Research Institute, University of Edinburgh, Edinburgh EH164TJ, United Kingdom

The Lin-11, Isl-1, and Mec-3 (LIM) homeodomain transcription factor Isl-1 has been reported to be involved in pituitary development in the early stages of mouse embryogenesis. Our recent studies have shown that Isl-1 is mainly located in the pituitary gonadotropes throughout pituitary development and persists to adulthood. We still do not know the physiological functions of Isl-1 expression and its related mechanisms in the pituitary gland. The aim of the present study was to examine the hypothesis that Isl-1 is involved in regulating pituitary gonadotropin hormone (FSH/LH) production by activating *FSH β* and *LH β* gene expressions. We have shown that Isl-1 activates *FSH β* and *LH β* subunit promoters and endogenous gene transcription in L β T2 cells. In addition, Isl-1 overexpression significantly increased FSH synthesis and secretion but not LH. The actions of Isl-1 were not observed when the homeodomain or LIM1 domains are mutated. This demonstrates that Isl-1 induction of *FSH β* and *LH β* is by both direct and indirect binding of Isl-1 to DNA sequences. Furthermore, Isl-1 expression level was up-regulated in L β T2 cells after exposure to GnRH, activin, and leptin. However, RNA interference-induced knockdown of Isl-1 significantly reduced the effect of leptin but did not obviously influence the stimulating effects of GnRH and activin on LH and FSH production. In conclusion, the results demonstrate that the LIM-homeodomain transcription factor Isl-1 functions to increase *FSH β /LH β* gene transcription, and mediates the effects of leptin on gonadotropin synthesis. (*Endocrinology* 151: 4787–4800, 2010)

The pituitary gonadotropins, FSH and LH, are heterodimeric glycoproteins composed of a common α -subunit (α -glycoprotein hormone α -subunit), and specific β -subunits (*FSH β* or *LH β*). In males and females, both these hormones are essential for gonadal function including folliculogenesis, spermatogenesis, and gonadal steroids and protein production in both sexes (1–3). Although the structures of these hormones are known, the mechanisms that regulate the synthesis and differential secretion of FSH and LH are not fully understood (4).

It is generally thought that the secretion of GnRH from the hypothalamus is key in regulating the gene expression of the gonadotropin β -subunits (5–7). GnRH increases hormonal production by binding to its transmembrane receptor. This binding increases intracellular calcium, which stimulates protein kinase C to activate nuclear proteins, including MAPK family members (8, 9). In addition to GnRH, the synthesis and secretion of gonadotropins are also regulated by the stimulating and inhibiting actions of estrogen and androgen (10) and peptides such as activin,

ISSN Print 0013-7227 ISSN Online 1945-7170
Printed in U.S.A.

Copyright © 2010 by The Endocrine Society
doi: 10.1210/en.2010-0208 Received February 19, 2010. Accepted July 13, 2010.
First Published Online August 11, 2010

Abbreviations: aa, Amino acids; ChIP, chromatin immunoprecipitation; D, postnatal day; E, embryonic day; Egr-1, early growth response protein 1; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; HD, homeodomain; HEK, human embryonic kidney; HOMO, homeodomain; HPG, hypothalamus-pituitary-gonadal; ICC, immunocytochemistry; IHC, immunohistochemistry; JAK, Janus tyrosine kinase; Lhx3, LIM class of homeodomain protein-3; LIM, Lin-11, Isl-1, and Mec-3; OB-Ra, short form of the leptin receptor; OB-Rb, long form of the leptin receptor; p, phosphorylated; Pitx, paired-like homeodomain transcription factor; SF-1, steroidogenic factor 1; STAT, signal transducer and activator of transcription.

inhibin (11, 12), leptin, and bone morphogenetic proteins (13, 14).

Leptin is a protein hormone synthesized and secreted mainly by adipocytes (14). In addition to its well-known regulating effects on appetite and metabolism, leptin is also vital in reproduction. Mutations in either the leptin (14) or the receptor (15) genes results in infertility. Leptin administration reverses reproductive abnormalities in leptin-deficient (*ob/ob*) mice (16) and accelerates sexual maturation and puberty in normal female (17) and transgenic mice (18). Although it is generally thought that the hypothalamus is a key site of action, the mechanism of leptin action in the reproductive system remains elusive. Leptin acts within the hypothalamus to stimulate GnRH release, which then triggers the subsequent release of FSH and LH, and stimulates the development and function of gonads (19). However, it was shown that leptin's action on the hypothalamus-pituitary-gonadal (HPG) axis appears at different levels of this axis (20, 21). The pituitary is a direct target of leptin action because leptin receptors are expressed in the pituitaries of many animals (22–26), and the cell types expressing leptin receptors include gonadotropes. In addition, leptin induces a bell-shaped dose-response curve of LH and FSH release from incubated anterior pituitaries in rats, pigs, and cattle (27–29), with almost the same effect on release of LH as GnRH (28, 29). These results suggest that leptin plays an additional role in regulating fertility by directly modulating pituitary LH and FSH synthesis and secretion. However, the pituitary intracellular pathways responsible for this direct action are still unknown. Studies have shown that leptin functions through Janus tyrosine kinase (JAK)-signal transducer and activator of transcription (STAT) signals in brain and other tissues (30–32).

The extrapituitary regulators, such as GnRH and activin, modulate LH β and FSH β subunit gene expression through a number of nuclear transcription factors. The factors involved in LH β transcription include steroidogenic factor 1 (SF-1), early growth response protein 1 (Egr-1), paired-like homeodomain transcription factor (Pitx)-1, p8, Sp1, and nuclear factor Y (33, 34). FSH β gene expression is mediated by SF-1, nuclear factor Y, activator protein-1, LIM class of homeodomain protein-3 (Lhx3), and Pitx class factors (34). The interplay among the different factors and genes are not fully understood.

The Lin-11, Isl-1, and Mec-3 (LIM)-homeodomain (HD) transcription factor Isl-1 is comprised of two tandem LIM domains and a HD. The homeodomain with a helix-turn-helix structure binds to regulatory DNA sequences of its target genes. The LIM domains are mainly involved in protein-protein interactions that regulate the activity of LIM-HD (35). Isl-1 was originally shown to function as an

insulin gene enhancer binding protein (36). It plays critical roles in cell determination, proliferation and differentiation in the nervous system (37, 38), heart (39), and pituitary gland (40). In mice, it is specifically expressed in the pituitary pouch rudiment at early development stages (41). *Isl-1*^{-/-} mice fetuses die at approximately embryonic day (E) 10 and the analysis at E9.5 demonstrates that the oral pouch is small and primitive with a conspicuously thinner wall (42). The exact function of Isl-1 on the fetal pituitary development is still unclear because *Isl-1* gene knockout causes developmental anomalies in the whole embryo (38), and pituitary cell differentiation happens mainly after E12.5 in the mouse (41). Our previous studies have shown that Isl-1 is mainly localized in gonadotropes, and Isl-1 expression parallels gonadotropic differentiation throughout the development in the sheep fetus and chicken embryo (43, 44). Additional studies should determine whether Isl-1 functions to promote pituitary gonadotrope differentiation and/or hormone production by activating FSH β and LH β genes.

The aims of the present study were to determine the effect of Isl-1 on LH β and FSH β gene expression and hormone production in gonadotropes and to identify whether Isl-1 is involved in mediating the signal pathways of GnRH, activin, and leptin in regulating LH and FSH production. The results show that Isl-1 enhances FSH β /LH β gene transcription and FSH production. It is out of our expectation that Isl-1 is involved in mediating the signal pathway of leptin but probably not GnRH and activin on gonadotropin production in L β T2 cells.

Materials and Methods

Animals and tissue collections

Kunming male mice were raised in a controlled temperature of 25 \pm 1 C on a 12-h light, 12-h dark cycle. The animal experiments were approved by the Chinese Association for Laboratory Animal Sciences. For immunohistochemistry, pituitary glands of three postnatal day (D) 40 mice were separated and fixed in 4% paraformaldehyde in PBS (pH 7.4) for 4 h at room temperature, embedded in paraffin, and cut into 5- μ m sections. In dietary manipulation experiments, 30 male mice at 21 D were randomly divided into three groups. Mice in the high-fat diet group were fed a diet containing 20% fat for 2 wk. The control and fasted groups were fed a regular grain diet, but the fasted group was starved for 48 h but with *ad libitum* drinking water. For determining serum leptin concentrations, the mice with dietary manipulation and normal mice aged 21 D (17 \pm 2 g), 40 D (24 \pm 2 g), and 60 D (37 \pm 3 g) were anesthetized, and blood samples were collected from the orbital sinus ($n \geq 6$ for each group). Serum was harvested and frozen at -20 C until RIA was conducted. After blood was taken, the mice were killed by cervical dislocation. Pituitaries were separated and divided into three factions per group for RT-PCR.

Immunohistochemistry (IHC) and immunocytochemistry (ICC)

Pituitary sections were dewaxed, rehydrated, and treated with 10% normal goat serum in PBS and incubated with rabbit antirat LH β (1:50; National Hormone and Peptide Program, Torrance, CA) and rabbit antihuman FSH β (1:50; Zymed, San Francisco, CA) at 4 C for 12 h. The section was then incubated with fluorescein isothiocyanate-labeled goat antirabbit IgG (1:50; Scottish Antibody Product Unit, Carlisle, UK) at room temperature for 3 h. After washing three times with PBS, the slides were observed under a fluorescence microscope (Leica Microsystems, Cambridge, UK) and photographed. Immunohistochemistry for Isl-1 detection was performed as previously described (43). The percentage of LH β or FSH β costaining for Isl-1 was counted.

For ICC, L β T2 cells were fixed in 4% paraformaldehyde for 10 min and permeabilized with 95% methanol at –20 C for 10 min and then blocked in 10% normal goat serum for 1 h. Subsequently, Isl-1 in the L β T2 cells was stained using the same protocol as IHC.

Plasmid construction

The –226 to +7 bp (–226LH β) and –712 to +7 bp (–712LH β) regions of the LH β gene and –1836 to +56 bp (–1836FSH β) region of FSH β gene were amplified from mouse genomic DNA by the PCR method using specific primers: LH- β , –226 bp, 5'-ACCTTGTTTCCCGTGCTT-3' and 5'-TCTTGAT-ACCTTCCCTACCTT-3'; LH- β , –712 bp, 5'-GACCGAATTT-GCCCAGTA-3' and 5'-TCTTGATACCTTCCCTACCTT-3'; FSH- β , –1836 bp, 5'-ATATAAACATCCCATCTCCA-3' and 5'-ATCAAGTGCTGCTACTCACC-3'. The –620 FSH β was obtained from the –1836 FSH β by *M*/fl restriction enzyme digestion. These fragments were respectively inserted into the pGL3.0-basic luciferase reporter vector (Promega, Madison, WI). The expression vector for murine Isl-1 pXJ40-myc-Isl-1 vector was kindly provided by Dr. Xinmin Cao (Institute of Molecular and Cell Biology, Singapore). The truncated mutants of Isl-1 were generated by restriction enzyme digestion and PCR methods. The LIM1 deletion (Δ Lim1) lacks amino acids (aa) 18–70, whereas the LIM2 deletion (Δ Lim2) lacks aa 95–109 in the Isl-1 peptide, and Δ Lim1 and -2 combines both deletions. The homeodomain mutated deletion (Δ Homo) lacks aa 193–213 in the Isl-1 homeodomain.

Cell culture and transient transfections

The mouse pituitary L β T2 and human embryonic kidney (HEK)-293T cells were cultured in 10% fetal bovine serum-DMEM (Life Technologies, Inc.-Invitrogen, Carlsbad, CA) supplemented with 100 IU/ml penicillin and 100 μ g/ml streptomycin and incubated at 37 C with 5% CO₂. Transient transfections or cotransfections were performed using Fugene HD or 6 transfection reagents (Roche Applied Science, Basel, Switzerland) according to the manufacturer's recommendations. All transfection experiments were performed at least three times.

Luciferase assays

L β T2 cells plated in 24-well plates were transfected with the Isl-1 expression vector or control vector, LH β or FSH β luciferase reporter vectors, and pTK-Ranilla vector (Promega) at a ratio of 10:4:1. Cells were harvested 24 h after transfection. Luciferase activity was measured using a dual-luc assay kit (Vigorous, Beijing,

China) on a Modulus microplate luminometer (Turner Biosystems, Sunnyvale, CA). The values shown by the fluc to rluc ratio were normalized to an empty luciferase reporter control.

HEK293T cells were cotransfected with pSilencer3.0-H1-Isl1ix (x indicates the targeted site) and pSilencer3.0-H1-shRNA control vector, respectively, with psiCHECK-Isl-1 (Promega) to estimate their interferential effects. Luciferase activity was measured 48 h later using a dual-luciferase reporter assay. Interference efficiency of each pSilencer3.0-H1-Isl1ix was shown by its rluc to fluc ratio to that of pSilencer3.0-H1-shRNA.

Radioimmunoassays

Transfected L β T2 cells (6×10^5 per 60 mm flask) were cultured in 10% fetal calf serum-DMEM for 24 h. Cell media were then replaced by DMEM (3 ml/flask). Cells were further incubated for 48 h before harvesting. In Isl-1 expression-interference experiments, cells were subjected to 10 ng/ml rat recombinant leptin (Abcam, Cambridge, UK), 10 nM GnRH analog (Sigma-Aldrich, St. Louis, MO), 25 ng/ml activin A (Peprotech, London, UK) or an equal volume DMEM for another 24 h. The media were then collected and the cells were lysed in 400 μ l lysis buffer for LH and FSH determinations. Cellular protein concentrations were determined by bicinchoninic assay. Experiments were performed four to six times. The hormone concentrations were normalized to the protein concentration.

LH, FSH, and leptin were analyzed using RIA reagents provided by the Beijing North Institute Biological Technology (Beijing, China). The minimum detectable concentrations were 1 mIU/ml for FSH and LH and 0.45 ng/ml for leptin. For each RIA the intra- and interassay coefficients of variation were less than 15% and less than 10%, respectively.

PCR and RT-PCR

Total RNA was extracted using Trizol, and cDNA was generated from 2 μ g RNA in a 25- μ l reaction mixture using Moloney murine leukemia virus (Promega) according to the manufacturer's protocols. LH β , FSH β , and internal control glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) gene expression were assayed by quantitative RT-PCR using primers as described previously (45–47). Isl-1 primers were designed by Primer Express 3.0 (Applied Biosystems, Foster City, CA) and confirmed to amplify a single product of the expected size via dissociation analysis and gel electrophoresis. Its sequences are: sense, 5'-AGT-CATCCGAGTGTGGTTTC-3', antisense, 5'-CATGCTGTT-GGGTGTATCTG-3'. Amplification was carried out on the ABI PRISM 7500 sequence detection system (Applied Biosystems), and each sample was assayed three times in duplicate. The relative abundance of the genes was determined using the ABI PRISM's software.

Leptin receptors, the short form (OB-Ra) and the long form (OB-Rb), were detected by PCR using primers as previous described (48). Amplifications were carried out on PCR instrument (Bio-Rad, Hercules, CA) using the following protocol: 94 C for 5 min (one time); 94 C for 50 sec; 54 C for 30 sec; 72 C for 30 sec (35 times); 72 C for 10 min; and 4 C holding.

Western blot analysis

Western blots were performed as described previously (49). Goat polyclonal Isl-1 antibody and mouse monoclonal GAPDH antibody were purchased, respectively, from R&D (Minneapolis, MN) and Ambion (Austin, TX). All experiments were performed three times. For quantification, the membranes were




FIG. 1. Isl-1 is expressed and associated with LH β and FSH β genes in the L β T2 gonadotropes. Dual labeling of LH β -Isl-1 (A–C) and FSH β -Isl-1 (D–F) were carried out in the anterior pituitary gland of mice. Green fluorescence shows the localization of LH β (B) and FSH β (E). Isl-1-positive cells in the same field of the same slide were stained brown (A and D). On their respective right are merged images (C and F). Purple arrows denote representative double-stained cells. Bar, 30 μ m. Three mice were examined in this experiment. A representative result is shown. G, ICC of Isl-1 in the L β T2 cells. Nuclear localizations of Isl-1 are stained dark brown. Bar, 30 μ m. H, Sequence analysis of mouse FSH β and LH β promoter regions of –1000 to +50 bp from the transcriptional sites. The putative Isl-1 binding sites are underlined. Arrows indicate the start site and the direction of transcription. Box sequences are primer binding sites of ChIP analysis. I, ChIP analysis. A cell aliquot before precipitation was designated as the input sample. PCR amplified 234-bp (LH β) or 401-bp (FSH β) regions in the promoters, respectively. Egr-1 was used as a positive control in precipitating the LH β promoter. IgG is a negative control provided by the kit.




FIG. 2. Is-1 induces the expressions of *LHB* and *FSHβ* genes and production of FSH. **A** and **B**, Effects of Is-1 expression vectors on the transiently transfected *LHB* (**A**) and *FSHβ* (**B**) gene promoters fused to luciferase reporter genes. The empty luciferase vector (luci) was also transfected as a control. The stimulating amount of Is-1-induced luciferase activity is denoted in parentheses. The experiments were repeated three times. **C–E**, Is-1 increased *LHB* and *FSHβ* mRNA levels. The pXJ40-myc-Is-1 and control vectors were transfected into LβT2 cells, respectively. Is-1 protein (**C**), *LHB* mRNA (**D**), and *FSHβ* mRNA (**E**) levels were analyzed 6, 15, and 24 h later by Western blot and RT-PCR, respectively. The present data of each assay were normalized to respective 6-h control (control vector transfected 6 h). Each sample was assayed three times in duplicate. **E** and **F**, Changes of LH (**E**) and FSH (**F**) levels after transfection of Is-1 expression vectors. The pXJ40-myc-Is-1 and control vectors were transfected into LβT2 cells, and concentration of LH and FSH in the media (secretion) and in the cellular extracts (synthesis) were determined 72 h later by RIA. In the data shown, hormone concentration is normalized to the corresponded protein concentration. Values are all means ± SEM from four to six separate experiments. *, $P < 0.05$; **, $P < 0.01$.

scanned on AlphaImager 2200 (Alpha Innotech Corp., San Leandro, CA). The results were analyzed by densitometry.

For ERK and STAT3 analysis, membranes were blocked and then incubated in 1:1000 diluted primary antibodies overnight at 4 C. The phosphorylated (p)-ERK1/2, p-STAT3, and total ERK1/2 antibodies were from Cell Signaling Technology (Beverly, MA). The STAT3 antibody was from Santa Cruz Biotechnology (Santa Cruz, CA). After incubation for 1–2 h with secondary antibodies (horseradish peroxidase conjugated), immunoreactive bands were visualized by chemiluminescence (Millipore, Bedford, MA). In each case, the membrane was initially probed with the antibody against the phosphorylated protein and then stripped using a buffer containing 2% sodium dodecyl sulfate, 62.5 mM Tris (pH 6.8), and 0.1 M 2-mercaptoethanol for 30 min at 50 C. After washing, immunodetection was repeated using an antibody against the total protein.

Chromatin immunoprecipitation (ChIP)

ChIP was performed using a ChIP assay kit (Active Motif, Carlsbad, CA) as previously described (50). Isl-1 antibody was from R&D. Antibodies of nonspecific IgG and Egr-1 (Santa Cruz Biotechnology) were used as negative and positive (only for LH β) controls, respectively. Some of the protein-DNA was not precipitated but set aside for total chromatin examination (termed input). After DNA was purified, the aimed sequence, from –226 to +7 of the mouse LH β gene and –687 to –286 bp of the FSH β gene, was amplified by PCR using the primer sequences of –226 LH β see *Plasmid construction*) and FSH- β : 5'-TACCATTGGCCCTA-AAT-3' and 5'-TTGGCGAACTCAGAATCCTT-3'.

Statistical analysis

All data were analyzed using one-way ANOVA, followed by Student's *t* test. All values are expressed as means \pm SEM. A *P* < 0.05 was considered significant.

Results

Isl-1 is expressed in the pituitary and gonadotrope-derived L β T2 cells

IHC results showed Isl-1 expression in the mouse pituitary gland. Dual-staining and cell-counting results showed that $64 \pm 5\%$ (mean \pm SEM) LH β immunopositive cells and $74 \pm 6\%$ (mean \pm SEM) of FSH β -immunopositive cells expressed Isl-1 in the mouse pituitary (Fig. 1, A–F). These data demonstrate that most gonadotropes express Isl-1 in the mouse pituitary. ICC results confirmed that Isl-1 was expressed in mouse gonadotrope-derived L β T2 cells (Fig. 1G).

Isl-1 interacts with LH β and FSH β genes in L β T2 cells

Sequence analysis showed that mouse FSH β and LH β genes contain two putative Isl-1 response elements (ATTAG) between –1000 and +50 bp upstream from the transcription sites (Fig. 1H). ChIP experiments were conducted to determine whether Isl-1 binds to the regulatory

regions of endogenous LH β and FSH β genes in L β T2 cells. The primers encompassing the ATTAG box were used to detect LH β and FSH β promoter sequences in genomic DNA precipitation (Fig. 1H). A 234-bp region of mouse LH β promoter was amplified from Isl-1-immunoprecipitated DNA as well as input chromatin and the DNA recruited by the LH β transcription factor Egr-1 (Fig. 1G, *top panel*). A 401-bp FSH β promoter sequence was also generated from Isl-1 immunoprecipitate (Fig. 1G, *bottom panel*). No PCR-amplified product was detected when the Isl-1 antibody was replaced by a nonspecific control antibody or in the water-template control (Fig. 1G). Taken together, these results confirm that Isl-1 interacts with the promoters of both FSH β and LH β in gonadotropes.




FIG. 3. Isl-1 mutants regulate –712LH β (A) and –620FSH β (B) promoter activities in L β T2 cells. The same quantity of wild and mutated Isl-1 expression vectors were cotransfected into cells with –712LH β or –620FSH β promoter reporter genes. Results are expressed as a ratio of firefly luciferase to renilla luciferase and then normalized to that of control. Δ Lim1, LIM1 domain mutated; Δ Lim2, LIM2 domain mutated; Δ Lim1 and 2, both LIM1 and LIM2 domain mutated; Δ Homo, homeodomain mutated. The experiments were repeated three times. Data are shown as means \pm SEM. Significant differences (*P* < 0.05) are shown by different letters.

Isl-1 increases FSH β and LH β promoter activity and affects hormone production

To determine whether Isl-1 is involved in regulating LH β and FSH β gene transcription, pXJ40-myc-Isl-1 was transfected into L β T2 cells with either LH β or FSH β promoter-luciferase genes. A dual-luciferase reporter assay showed that Isl-1 enhanced the promoter activities of –226LH β and –712LH β about 2.5-fold ($P < 0.05$) and 2.3-fold ($P < 0.01$), respectively (Fig. 2A). –620FSH β and –1836FSH β were also stimulated by Isl-1 about 6.1-fold ($P < 0.01$) and 1.9-fold ($P < 0.05$), respectively (Fig. 2B). Meanwhile, Isl-1 overexpression did not change the luciferase activity of pGL3.0-basic ($P > 0.05$).

To determine whether the overexpression of Isl-1 affects endogenous transcriptional activity of gonadotropin β -subunits, we examined variations of their mRNA levels in pXJ40-myc-Isl-1- or control vector-transfected L β T2 cells. Western blot results indicated that Isl-1 expression did not increase until 15 and 24 h after transfection of pXJ40-myc-Isl-1 (Fig. 2C). Correspondingly, both LH β and FSH β transcripts were significantly elevated to about 1.5-fold at 15 and 24 h ($P < 0.01$) but were not changed at 6 h ($P > 0.05$, Fig. 2, D and E). In contrast, there was no change in LH β or FSH β transcripts in control vector-transfected cells throughout ($P > 0.05$, Fig. 2, D and E).

Although Isl-1 significantly increased both LH β and FSH β mRNA levels, RIA results showed that Isl-1 resulted in a significant increase only in intracellular ($P < 0.01$) and released ($P < 0.05$) FSH (Fig. 2G) but not LH ($P > 0.05$, Fig. 2F).

Isl-1 HOMO and LIM domains regulate FSH β and LH β promoters' activities

LIM-HD Isl-1 has one homeodomain and two separated LIM domains, interacting with target DNAs and proteins, respectively. To determine which domains are involved in the observed regulation of LH β and FSH β

gene transcription in L β T2 cells, four truncated forms of Isl-1 protein were constructed: Δ Lim1, Δ Lim2, Δ Lim1 and 2, and Δ Homo. Activation of Isl-1 on both –712LH β and –620FSH β was decreased ($P < 0.05$, Fig. 3) when LIM1, LIM1 and 2, or the homeodomain was mutated. In contrast, deletion of LIM2 alone did not significantly change the stimulating effects of Isl-1, whether on –712LH β or –620FSH β ($P > 0.05$, Fig. 3).

GnRH, activin, and leptin all increase the expression levels of LIM-HD Isl-1

To determine the potential for Isl-1 to mediate the effects of GnRH, activin, and leptin on gonadotropins, we determined mRNA levels of Isl-1 in L β T2 cells exposed to GnRH, activin, and leptin using RT-PCR. After GnRH treatment, there was a significant increase in the Isl-1 mRNA level at 1 h ($P < 0.01$), which peaked at 3 h ($P < 0.01$) and decreased thereafter to the control level by 24 h ($P > 0.05$, Fig. 4A). In response to activin and leptin, Isl-1 mRNA level increased by 3 h ($P < 0.05$), peaked at 12 h, and stayed for at least 24 h ($P < 0.01$, Fig. 4, B and C). These results suggest that the LIM-HD transcription factor Isl-1 is potentially involved in gonadotropin-regulating effects of GnRH, activin, and leptin in gonadotropes.

Knockdown of Isl-1 expression in the L β T2 cells

To directly test the role of Isl-1, five short hairpin RNA constructs targeting mouse *Isl-1* gene (Fig. 5A) were designed to inhibit Isl-1 expression in L β T2 cells. Each construct was screened individually for inhibiting the activation of the psiCHECK-Isl-1 luciferase in transient transfection assays. pSilencer-H1-Isl1891 was the best inhibitor, reducing Isl-1 expression by about 75%, whereas Isl1511 resulted in a 64% reduction. Isl1521, Isl1138, and Isl1288 resulted in a 49, 35, and 37% reduction, respectively (Fig. 5B).




FIG. 4. The relative levels of Isl-1 mRNA levels in L β T2 cells treated by GnRH (A), activin (B), and leptin (C) for different times. Cells were treated with 10 nM GnRH, 25 ng/ml activin, or 10 ng/ml leptin for 3, 6, 12, and 24 h. mRNA levels of Isl-1 and reference gene, *GAPDH*, were determined by RT-PCR. Data are shown as means \pm SEM from three experiments in duplicate and normalized to their respective control. *, $P < 0.05$; **, $P < 0.01$.




FIG. 5. Silencing effects of Isl-1 interference vector. A, Sketch map indicates the targeted sites of several Isl-1 interference vectors (named pSilencer-H1-Islix) in the mouse Isl-1 cDNA. B, psiCHECK-Isl-1 was transfected into HEK293T cells together with pSilencer-H1-Islix. Dual-luciferase ratios were expressed as ratios of renilla luciferase to firefly luciferase, and they were normalized to pSilencer-H1-shRNA control. Data are shown as means \pm SEM from three repeated experiments. *, $P < 0.05$; **, $P < 0.01$. C and D, Western blot analysis of Isl-1 protein levels in L β T2 cells transfected with pSilencer-H1-shRNA, pSilencer-H1-Isl1891, and nontreated cells (NT). GAPDH is blotted as an internal control. The integrated density value (IDV) of Isl-1 and GAPDH products were determined using the Alphamager 2200 software package (Alpha Innotech), and Isl-1 expression levels were expressed according to IDV. Data from three experiments were all normalized to the control and are shown as means \pm SEM. Significant differences ($P < 0.05$) are shown by different letters.

To assess the pSilencer-H1-Isl1891-inhibiting effects in gonadotropes, protein levels of Isl-1 were determined in L β T2 cells. Isl-1 protein levels in empty vectors and nontreated cells did not change significantly ($P > 0.05$) but decreased about 40.2% ($P < 0.05$, Fig. 5, C and D) in cells transfected with pSilencer-H1-Isl1891, confirming its efficacy of reducing Isl-1 expression.

Isl-1 is involved in leptin's regulation of LH and FSH synthesis

The intracellular and medium concentrations of LH and FSH in Isl-1 knockdown and control cells were determined after treatment with GnRH, activin, and leptin to assess the role of Isl-1 in their functions.

In Isl-1 knockdown cells in the absence of GnRH, activin, or leptin (Fig. 6, NT group), the levels of secreted FSH decreased ($P < 0.05$, Fig. 6B). There was no effect on intracellular or secreted LH levels ($P > 0.05$).

GnRH treatment caused a small increase of LH secretion in L β T2 cells ($P < 0.05$), and Isl-1 knockdown did not influence this change ($P > 0.05$, Fig. 6A). Activin did not change the levels of LH ($P > 0.05$, Fig. 6A); however, it significantly promoted both synthesized ($P < 0.05$) and secreted ($P < 0.01$) FSH levels in wild-type cells, which also were not changed by Isl-1 interference ($P > 0.05$, Fig. 6B). In contrast, leptin increased intracellular LH ($P < 0.05$, Fig. 6A) and FSH ($P < 0.05$, Fig. 6B) concentrations in control vector-transfected cells. Both were significantly ($P < 0.01$ for LH, Fig. 6A; $P < 0.05$ for FSH, Fig. 6B) decreased in Isl-1 knockdown cells, indicating that Isl-1 mediates leptin regulation of LH and FSH synthesis.

Leptin induces Isl-1 expression by JAK-STAT

PCR results showed that both the OB-Ra and OB-Rb forms of the leptin receptor are expressed in L β T2 cells as well as in mouse pituitary, hypothalamus, and ovary (Fig. 7A). We next examined leptin-stimulated signals in L β T2 cells. Cultured cells were exposed to leptin for up to 60 min. Cell lysates were immunoprecipitated with antiphospho-STAT3 or antiphospho-ERK antibodies. The leptin-dependent increases in STAT3 and ERK activity were both detectable at 5 min (Fig. 7B), followed by a sustained increase from 5 to 60 min (Fig. 7B). These results indicate that leptin activates both STAT3 and ERK signals in L β T2 cells. In addition, leptin-induced expression of Isl-1 mRNA was blocked by both the JAK-selective tyrosine kinase inhibitor AG490 and STATs' phosphorylation suppressor parthenolide ($P < 0.05$, Fig. 7D). In contrast, PD98059, a specific antagonist of ERK1/2, did not modify the transcription level of Isl-1 induced by leptin ($P > 0.05$). Interestingly, PD98059 alone also increased basal Isl-1 expressions ($P < 0.05$, Fig. 7D).

The relation of pituitary Isl-1 mRNA levels to serum leptin concentration

To further identify the potential relationship of Isl-1 and leptin *in vivo*, we measured the pituitary Isl-1 mRNA by RT-PCR and serum leptin levels by RIA in mice at different ages and with different feeding patterns. The results showed that Isl-1 mRNA increased significantly from




FIG. 6. Isl-1 is involved in leptin regulation of gonadotropin synthesis. pSilencer-H1-Isl-1891 and control vectors were transfected into L β T2 cells, and after 48 h, 10 nM GnRH, 25 ng/ml activin and 10 ng/ml leptin were respectively added 24 h before harvest. The LH (A) and FSH (B) hormone concentration in media and cellular extracts were determined by RIA, and the data are shown by hormone concentration normalized to the corresponding cellular protein concentration. Values are means \pm SEM from four to six separate experiments. Significant differences are indicated by different letters ($P < 0.05$). NT, Nontreated cells.

21 D to 60 D ($P < 0.05$, Fig. 8A), whereas serum leptin levels increased significantly only at 60 D ($P < 0.05$, Fig. 8B). However, the changes of pituitary Isl-1 mRNA level did parallel global leptin concentrations in dietary manipulated mice, with both increasing significantly ($P < 0.01$) after feeding with the high-fat diet but were not significantly ($P > 0.05$) changed by starvation for 48 h (Fig. 8, C and D).

Discussion

Isl-1 expression in the developing pituitary gland has been reported in mice (41, 51), chicks (44, 52), and sheep (43). The dominant cell types expressing Isl-1 in the pituitary gland are the FSH β - and LH β -expressing gonadotropes. Our present results agree with these findings and show that Isl-1 is mainly expressed in pituitary gonadotropes of prepubertal to adult mice and in gonadotrope-derived cell line, L β T2. The discovery that L β T2 cells express Isl-1 provides an *in vitro* tool for studying the transcriptional effect of Isl-1 on LH/FSH β gene expression and hormone production.

Previous studies on the function of Isl-1 in the pituitary have focused on the early stage of embryo development.

Isl-1 null mice embryos show developmental anomalies at E9.5 and die at E10 (38, 42), 2.5 d before pituitary cell differentiation and hormone production (41). Therefore, this Isl-1 gene-knockout animal model cannot be used to study the function of Isl-1 in pituitary development and hormone secretion. Upon that premise, the present study switches to *in vitro* study, to demonstrate the relationship of Isl-1 and gonadotropin.

First, we have shown that Isl-1 not only increases both LH β and FSH β promoter activities but also elevates their mRNA levels about 1.5-fold. These changes in mRNA agree with luciferase assay results, which indicated that Isl-1 causes a similar stimulation on long-form LH β and FSH β promoter genes, which are, respectively, increased 2.3- and 1.9-fold. The elevation of mRNA strongly indicates that Isl-1 probably affects synthesis and secretion of gonadotropins by altering LH β and FSH β gene expression, which should result in increased hormone production. However, the results have shown that both synthesis and secretion levels of FSH are increased by Isl-1 in L β T2 cells but not that of LH.

Differential responses of LH and FSH may relate to their different synthesis and secretion patterns. Indeed, the LH and FSH β subunits have quite distinct amino acid sequences and variable oligosaccharide chains (53). Glycosylation of polypeptides is a posttranslational event, which influences the rate of hormone dimer assembly (54) and thus synthesis. Furthermore, LH and FSH also exhibit different secretion patterns. Previous studies in sheep have confirmed that LH is stored in electron-dense granules within the gonadotrope (55) associated with SgII (56–58), and FSH appears in electron light granules, possibly associated with chromogranin A (57–59). The release and subsequent plasma concentrations of FSH appear more closely related to the amount of FSH being produced within the gonadotrope, and also to the levels of transcription of FSH β mRNA (4). In contrast the release of LH occurs via a regulated pathway with little relationship to the levels of transcription of LH β mRNA. This is probably why LH was not changed by Isl-1, although the expression level of LH β mRNA was up-regulated.

Similar to other LIM-HD transcription factors, Isl-1 contains two LIM domains and a HD. To study the roles of these domains in regulating LH β and FSH β gene expression, we constructed four Isl-1 mutants. The results showed that the HD is required for full Isl-1 transcrip-




FIG. 7. Intracellular signals of leptin in up-regulating Isl-1 in the L β T2 cells. **A**, Expressions of leptin receptor mRNAs in L β T2 cells and several tissues of the adult mouse. Lane 1, DNA ladder; lane 2, mouse hypothalamus; lane 3, mouse pituitary; lane 4, L β T2 cell; lane 5, mouse ovary; lane 6, reverse transcriptase blank without Moloney murine leukemia virus. OB-Ra and OB-Rb products are, respectively, 237 and 446 bp. GAPDH is used as a positive control. Experiments were repeated three times, and one representative is shown. **B**, Leptin activates ERK and STAT3 signals in the L β T2 cells. The L β T2 cells were cultured with 10 ng/ml leptin for 0 (no treatment control), 5, 10, 20, 30, and 60 min, respectively, and then total protein extracts were prepared. Levels of phosphorylated ERK, phosphorylated STAT3, total ERK, and total STAT3 proteins were detected by Western blot. **C**, Changes of p-STAT3/STAT3 and p-ERK/ERK Levels. Data were calculated using integrated density value ratio of bands in **B**, which were determined by Alphamager 2200 (Alpha Innotech). **D**, The relative levels of Isl-1 mRNA in L β T2 cells treated by leptin and several inhibitors. Inhibitors were added 1 h before leptin. All inhibitors had a final concentration of 20 μ M. After 12 h exposure to leptin, Isl-1 expressions in each group were determined by RT-PCR and then normalized to that of control. Data are shown as means \pm SEM from three repeated experiments. Significance ($P < 0.05$) is shown by different letters. NT, Nontreated cells; PTL, parthenolide.

tional activity on LH β and FSH β genes, suggesting that Isl-1 directly interacts with LH β and FSH β DNA. ChIP results confirmed that the binding site of Isl-1 probably lies within the regions of -226 to $+7$ bp on the LH β genes and -687 to -286 bp on the FSH β gene, although the exact binding motifs requires further work. In addition, the full transcriptional activity of Isl-1 does not occur when the LIM1 domain is absent. This indicates that Isl-1 also influences LH β and FSH β transcription by interacting with other proteins through LIM1. Isl-1 is capable of interacting with Lhx3 and the LIM-interacting partners, Ldb1, to form a trimer using its ligand-binding domain and LIM1 domains (60). Isl-1 also functions in synergy with other transcription factors, such as SF-1 (51), hepatocyte nuclear factor-4 α (61), and class B bHLH transcription factor, BETA2 (62). The specific associating proteins were not identified in this study. Possibly, one of the reasons of

Isl-1's different magnitude of the effect on the LH β and FSH β promoter activities is related to different specific transcription factor interactions, *e.g.* Lhx3 has been shown to regulate FSH β but not LH β promoters (63). Collectively, it is likely that the Isl-1-mediated regulation of the LH β and FSH β promoters incorporates several mechanisms involving both direct and indirect interactions.

More importantly, the regulatory effect on FSH synthesis and secretion potentially extends the role of Isl-1 to the HPG axis. GnRH, steroids, and cytokines produced by the hypothalamus and gonads possibly regulate the expression or activation of Isl-1 in the pituitary, thus increasing FSH hormone production. Hormone measurements demonstrate that GnRH induces LH but not FSH secretion, and activin stimulates FSH but not LH secretion, which is in agreement with previous studies (7, 45, 64). However, Isl-1 does not appear to have a close rela-




FIG. 8. Changes of pituitary Isl-1 mRNA levels and serum leptin by age and diet. A and B, Pituitary Isl-1 mRNA levels and serum leptin concentrations in mice 21, 40, and 60 d old. Isl-1 mRNA levels were determined by real-time PCR. GAPDH is a reference gene. Each sample was assayed three times in duplicate. Present data of each assay was normalized to that of the 21 D group. The sample numbers for leptin determination were $n = 6$ (21 D), $n = 6$ (40 D), and $n = 10$ (60 D), respectively. C and D, Changes of pituitary Isl-1 mRNA levels and serum leptin concentrations after 2 wk of dietary manipulation. The control group was fed a regular grain diet; the high-fat diet (HFD) group was fed a diet containing 20% fat. Mice in the fast group were starved for the last 48 h with sufficient drinking water. The sample number for each group is 10. Isl-1 mRNA levels were determined by real-time PCR. Each sample was assayed three times in duplicate and normalized to that of control group. All of the data are shown as means \pm SEM. *, $P < 0.05$; **, means $P < 0.01$.

tionship with the function of GnRH and activin because their regulatory effects on gonadotropin are not changed by Isl-1 interference. This is reasonable because GnRH has been shown to directly regulate *LHB* through *Egr-1*, which synergizes with SF-1 and Pitx1 (65), and activin regulates FSH synthesis by small mothers against decapentaplegic homolog (smad) 2/3, which acts directly on the *FSH* promoter (66). However, GnRH and activin increase Isl-1 expression, and the effects of GnRH and activin on FSH and LH do not rely on Isl-1. It is possible that Isl-1 may be involved in other GnRH- and activin-induced effects, e.g. cell proliferation, as shown in other cells or tissues (67, 68). This requires further studies.

The results of the present study show that leptin stimulates both FSH and LH synthesis in L β T2 cells, in agreement with some previous reports (27, 28, 69). However, there are conflicting reports about the effect of leptin on gonadotropins and the HPG axis. Administration of leptin *in vitro* reduced LH and FSH secretion in the rat hemipituitary cultures (70) and failed to stimulate FSH in energy-deprived rhesus monkeys (71). Furthermore, in the pres-

ence of sufficient energy stores, leptin did not alter reproductive function (19). It is unclear why these discrepancies in the effect of leptin occur. It may be due to the differential effects of leptin in modulating the expression of its own receptors. It may also be due to the dose dependence of leptin. Indeed, the concentration of endogenous leptin is at a comparatively high level in the animals with sufficient energy stores, and this has been confirmed in our results. Thus, leptin may already be acting at maximum. An overdose of leptin was demonstrated to be significantly decreased the expressional levels of hypothalamic OB-Rb (72). Furthermore, there is evidence that leptin's lack of effectiveness on the HPG axis may be due to influences of other neuroendocrine peptides, like neuropeptide Y, GnRH, and estrogen (73–75).

In the reproductive axis, leptin is known to activate signaling via its long- and short-form receptors. And we have now shown that both forms of the receptor are expressed in the hypothalamus, pituitary, and ovary in the mouse, confirming a previous report (22). Studies in cattle (23) and sheep (24) pituitaries have suggested that most of

OB-R-positive cells are gonadotropes. This agrees well with present result that OB-Rs are expressed in L β T2 gonadotropes.

The present work is the first to show, using specific inhibitors, that Isl-1 mRNA levels are regulated by leptin through JAK-STAT's cascade in pituitary gonadotropes, similar to the hypothalamus (76, 77). This, however, differs with a previous report that shows that leptin's action in the reproductive neuroendocrine axis is independent of pSTAT3 signaling as shown by disrupting Tyr¹¹³⁸ phosphorylation in the OB-Rb (78). It has been reported that JAK-stimulated Isl-1 could induce Tyr phosphorylation, DNA binding activity, and target gene expression of STAT3 independent of other activators (e.g. epithelial growth factor) (79). This suggests that Isl-1 may function as an adaptor protein that brings JAK and STAT3 into proximity and thereby facilitates STAT3 phosphorylation by JAK. This fits well with our results, which indicate that leptin elevates Isl-1 expression, resulting in the induction of STAT3 phosphorylation. However, the inhibitor parthenolide used in our study is a broad inhibitor of phosphorylation of STATs, including STAT1, -3, -5, and -6. We did not check the phosphorylation of other STATs, which may have roles in the present study. This study also shows that leptin activates the ERK pathway in L β T2 gonadotrope cells. It is in agreement with the report that leptin promotes proliferation via the MAPK/ERK signal pathway in endometrial cancer cells (80). This suggests that leptin may function to promote L β T2 cell proliferation by activating ERK. Nevertheless, the molecular mechanism of leptin's effect through STATs and MAPK need to be elucidated further.

In the present study, serum leptin rose from 40 D to 60 D, which agrees with a previous study that showed leptin increases from 20 D to postpuberty in male mouse plasma (81). The expression levels of pituitary Isl-1 also increased over this time. This parallel change suggests that leptin is an important determinative regulator for Isl-1 around puberty. However, the changes in leptin and Isl-1 were not completely consistent with serum LH and FSH levels (82). This is probably related to the complex effects of other regulators like activin, inhibin, and principally GnRH during the prepubertal-pubertal transition. More importantly, our results also show that serum leptin and pituitary Isl-1 display parallel changes under dietary manipulations. This, together with the developmental results, strongly supports the *in vitro* results demonstrating that leptin regulates pituitary Isl-1 expression and reveals their close physiological relationship *in vivo*.

In conclusion, this novel study demonstrates that Isl-1 not only increases the expression of LH and FSH β -sub-

unit genes but also mediates the regulating functions of leptin on LH and FSH synthesis.

Acknowledgments

We thank Professor Pamela L. Mellon (University of California, San Diego, San Diego, CA) for providing the L β T2 cell lines. pXJ40-myc-Isl-1 plasmid was a gift of Dr. Xinmin Cao (Institute of Molecular and Cell Biology, Singapore). The 40.2D6 Isl-1 antibody was developed by Thomas Jessell and obtained from the Developmental Studies Hybridoma Bank (Iowa City, IA) developed under the auspices of the National Institute of Child Health and Child Development.

Address all correspondence and requests for reprints to: Sheng Cui, State Key Laboratory of Agrobiotechnology, College of Biological Sciences, China Agricultural University, Beijing 100193, People's Republic of China. E-mail: cuisheng@cau.edu.cn.

This work was supported by the Natural Science Foundation of China (Grants 30630011 and 30771587), the National Basic Research Program of China (Grants 2007CB947402 and 2009CB941702), and the Innovative Project of State Key Laboratory of Agrobiotechnology (Grant 2009SKLAB05-4). A.S.M. was supported by the United Kingdom Medical Research Council Grant G7.00007.01.

Disclosure Summary: The authors have nothing to disclose.

References

1. Ma X, Dong Y, Matzuk MM, Kumar TR 2004 Targeted disruption of luteinizing hormone β -subunit leads to hypogonadism, defects in gonadal steroidogenesis, and infertility. *Proc Natl Acad Sci USA* 101:17294–17299
2. Vegetti W, Alagna F 2006 FSH and folliculogenesis: from physiology to ovarian stimulation. *Reprod Biomed Online* 12:684–694
3. Allan CM, Handelsman DJ 2005 Transgenic models for exploring gonadotropin biology in the male. *Endocrine* 26:235–239
4. McNeilly AS, Crawford JL, Taragnat C, Nicol L, McNeilly JR 2003 The differential secretion of FSH and LH: regulation through genes, feedback and packaging. *Reprod Suppl* 61:463–476
5. Millar RP, Pawson AJ, Morgan K, Rissman EF, Lu ZL 2008 Diversity of actions of GnRHs mediated by ligand-induced selective signaling. *Front Neuroendocrinol* 29:17–35
6. Burger LL, Haisenleder DJ, Dalkin AC, Marshall JC 2004 Regulation of gonadotropin subunit gene transcription. *J Mol Endocrinol* 33:559–584
7. Miller WL, Shafiee-Kermani F, Strahl BD, Huang HJ 2002 The nature of FSH induction by GnRH. *Trends Endocrinol Metab* 13:257–263
8. Caunt CJ, Finch AR, Sedgley KR, McArdle CA 2006 GnRH receptor signalling to ERK: kinetics and compartmentalization. *Trends Endocrinol Metab* 17:308–313
9. Dobkin-Bekman M, Naidich M, Pawson AJ, Millar RP, Seger R, Naor Z 2006 Activation of mitogen-activated protein kinase (MAPK) by GnRH is cell-context dependent. *Mol Cell Endocrinol* 252:184–190
10. Thackray VG, Mellon PL 2008 Synergistic induction of follicle-stimulating hormone β -subunit gene expression by gonadal steroid

- hormone receptors and Smad proteins. *Endocrinology* 149:1091–1102
11. Padmanabhan V, McNeilly AS 2001 Is there an FSH-releasing factor? *Reproduction* 121:21–30
 12. Risbridger GP, Schmitt JF, Robertson DM 2001 Activins and inhibins in endocrine and other tumors. *Endocr Rev* 22:836–858
 13. Otsuka F, Shimasaki S 2002 A novel function of bone morphogenetic protein-15 in the pituitary: selective synthesis and secretion of FSH by gonadotropes. *Endocrinology* 143:4938–4941
 14. Strobel A, Issad T, Camoin L, Ozata M, Strosberg AD 1998 A leptin missense mutation associated with hypogonadism and morbid obesity. *Nat Genet* 18:213–215
 15. Clément K, Vaisse C, Lahlou N, Cabrol S, Pelloux V, Cassuto D, Goumelen M, Dina C, Chambaz J, Lacorte JM, Basdevant A, Bougnères P, Lebouc Y, Froguel P, Guy-Grand B 1998 A mutation in the human leptin receptor gene causes obesity and pituitary dysfunction. *Nature* 392:398–401
 16. Chehab FF, Lim ME, Lu R 1996 Correction of the sterility defect in homozygous obese female mice by treatment with the human recombinant leptin. *Nat Genet* 12:318–320
 17. Chehab FF, Mounzih K, Lu R, Lim ME 1997 Early onset of reproductive function in normal female mice treated with leptin. *Science* 275:88–90
 18. Yura S, Ogawa Y, Sagawa N, Masuzaki H, Itoh H, Ebihara K, Aizawa-Abe M, Fujii S, Nakao K 2000 Accelerated puberty and late-onset hypothalamic hypogonadism in female transgenic skinny mice. *J Clin Invest* 105:749–755
 19. Cunningham MJ, Clifton DK, Steiner RA 1999 Leptin's actions on the reproductive axis: perspectives and mechanisms. *Biol Reprod* 60:216–222
 20. Foster DL, Nagatani S 1999 Physiological perspectives on leptin as a regulator of reproduction: role in timing puberty. *Biol Reprod* 60:205–215
 21. Herrid M, Xia Y, O'Shea T, McFarlane JR 2008 Leptin inhibits basal but not gonadotrophin-stimulated testosterone production in the immature mouse and sheep testis. *Reprod Fertil Dev* 20:519–528
 22. Zamorano PL, Mahesh VB, De Sevilla LM, Chorich LP, Bhat GK, Brann DW 1997 Expression and localization of the leptin receptor in endocrine and neuroendocrine tissues of the rat. *Neuroendocrinology* 65:223–228
 23. Ogasawara H, Ohwada S, Nagai Y, Taketa Y, Matsuzaki M, Tanaka S, Watanabe K, Aso H, Yamaguchi T 2008 Localization of leptin and leptin receptor in the bovine adenohypophysis. *Domest Anim Endocrinol* 35:8–15
 24. Iqbal J, Pompolo S, Considine RV, Clarke IJ 2000 Localization of leptin receptor-like immunoreactivity in the corticotropes, somatotropes, and gonadotropes in the ovine anterior pituitary. *Endocrinology* 141:1515–1520
 25. Jin L, Zhang S, Burguera BG, Couce ME, Osamura RY, Kulig E, Lloyd RV 2000 Leptin and leptin receptor expression in rat and mouse pituitary cells. *Endocrinology* 141:333–339
 26. Lin J, Barb CR, Matteri RL, Kraeling RR, Chen X, Meinersmann RJ, Rampacek GB 2000 Long form leptin receptor mRNA expression in the brain, pituitary, and other tissues in the pig. *Domest Anim Endocrinol* 19:53–61
 27. Barb CR, Kraeling RR 2004 Role of leptin in the regulation of gonadotropin secretion in farm animals. *Anim Reprod Sci* 82–83:155–167
 28. Yu WH, Kimura M, Walczewska A, Karanth S, McCann SM 1997 Role of leptin in hypothalamic-pituitary function. *Proc Natl Acad Sci USA* 94:1023–1028
 29. Tezuka M, Irahara M, Ogura K, Kiyokawa M, Tamura T, Matsuzaki T, Yasui T, Aono T 2002 Effects of leptin on gonadotropin secretion in juvenile female rat pituitary cells. *Eur J Endocrinol* 146:261–266
 30. Frühbeck G 2006 Intracellular signalling pathways activated by leptin. *Biochem J* 393(Pt 1):7–20
 31. Cirillo D, Rachiglio AM, la Montagna R, Giordano A, Normanno N 2008 Leptin signaling in breast cancer: an overview. *J Cell Biochem* 105:956–964
 32. Hegyi K, Fülöp K, Kovács K, Tóth S, Falus A 2004 Leptin-induced signal transduction pathways. *Cell Biol Int* 28:159–169
 33. Brown P, McNeilly AS 1999 Transcriptional regulation of pituitary gonadotropin subunit genes. *Rev Reprod* 4:117–124
 34. Melamed P, Kadir MN, Wijeweera A, Seah S 2006 Transcription of gonadotropin β subunit genes involves cross-talk between the transcription factors and co-regulators that mediate actions of the regulatory hormones. *Mol Cell Endocrinol* 252:167–183
 35. Curtiss J, Heilig JS 1998 DeLIMiting development. *Bioessays* 20:58–69
 36. Karlsson O, Thor S, Norberg T, Ohlsson H, Edlund T 1990 Insulin gene enhancer binding protein Isl-1 is a member of a novel class of proteins containing both a homeo- and a Cys-His domain. *Nature* 344:879–882
 37. Ericson J, Thor S, Edlund T, Jessell TM, Yamada T 1992 Early stages of motor neuron differentiation revealed by expression of homeobox gene *Islet-1*. *Science* 256:1555–1560
 38. Pfaff SL, Mendelsohn M, Stewart CL, Edlund T, Jessell TM 1996 Requirement for LIM homeobox gene *Isl-1* in motor neuron generation reveals a motor neuron-dependent step in interneuron differentiation. *Cell* 84:309–320
 39. Cai CL, Liang X, Shi Y, Chu PH, Pfaff SL, Chen J, Evans S 2003 *Isl-1* identifies a cardiac progenitor population that proliferates prior to differentiation and contributes a majority of cells to the heart. *Dev Cell* 5:877–889
 40. Sheng HZ, Moriyama K, Yamashita T, Li H, Potter SS, Mahon KA, Westphal H 1997 Multistep control of pituitary organogenesis. *Science* 278:1809–1812
 41. Ericson J, Norlin S, Jessell TM, Edlund T 1998 Integrated FGF and BMP signaling controls the progression of progenitor cell differentiation and the emergence of pattern in the embryonic anterior pituitary. *Development* 125:1005–1015
 42. Takuma N, Sheng HZ, Furuta Y, Ward JM, Sharma K, Hogan BL, Pfaff SL, Westphal H, Kimura S, Mahon KA 1998 Formation of Rathke's pouch requires dual induction from the diencephalon. *Development* 125:4835–4840
 43. Liu J, Liu Z, Yi S, Cui S 2005 *Islet-1* expression and its colocalization with luteinizing hormone, thyroid-stimulating hormone and oestrogen receptor α in the developing pituitary gland of the sheep foetus. *J Neuroendocrinol* 17:773–780
 44. Liu J, He Y, Wang X, Zheng X, Cui S 2005 Developmental changes of *Islet-1* and its co-localization with pituitary hormones in the pituitary gland of chick embryo by immunohistochemistry. *Cell Tissue Res* 322:279–287
 45. Nicol L, McNeilly JR, Stridsberg M, McNeilly AS 2004 Differential secretion of gonadotropins: investigation of the role of secretogranin II and chromogranin A in the release of LH and FSH in *L β T2* cells. *J Mol Endocrinol* 32:467–480
 46. Su P, Wu JC, Sommer JR, Gore AJ, Petters RM, Miller WL 2005 Conditional induction of ovulation in mice. *Biol Reprod* 73:681–687
 47. Uehara T, Kaneko M, Tanaka S, Okuma Y, Nomura Y 1999 Possible involvement of p38 MAP kinase in HSP70 expression induced by hypoxia in rat primary astrocytes. *Brain Res* 823:226–230
 48. Frühbeck G, Gómez-Ambrosi J, Martínez JA 1999 Pre- and postprandial expression of the leptin receptor splice variants OB-Ra and OB-Rb in murine peripheral tissues. *Physiol Res* 48:189–195
 49. Wang G, Chen D, Luo H, Liu J, Ji X, Fan J, Cui S 2010 Low-dose ethanol suppresses 17 β -estradiol activity in GH4C1 pituitary tumor cells. *Cell Biol Toxicol* 26:265–277
 50. Liu S, Luo H, Liu J, McNeilly AS, Cui S 2008 Specificity protein 1 (Sp1) plays role in regulating LIM homeodomain transcription factor *Lhx4* gene expression. *Biochem Biophys Res Commun* 366:36–41
 51. Granger A, Bleux C, Kottler ML, Rhodes SJ, Counis R, Laverrière JN 2006 The LIM-homeodomain proteins *Isl-1* and *Lhx3* act with

- steroidogenic factor 1 to enhance gonadotroph-specific activity of the gonadotropin-releasing hormone receptor gene promoter. *Mol Endocrinol* 20:2093–2108
52. Sjödal M, Gunhaga L 2008 Expression patterns of Shh, Ptc2, Raldh3, Pitx2, Isl-1, LIM3 and Pax6 in the developing chick hypothalamic placode and Rathke's pouch. *Gene Expr Patterns* 8:481–485
53. Jablonka-Shariff A, Boime I 2004 Luteinizing hormone and follicle-stimulating hormone exhibit different secretion patterns from cultured Madin-Darby canine kidney cells. *Biol Reprod* 70:649–655
54. Corless CL, Matzuk MM, Ramabhadran TV, Krichevsky A, Boime I 1987 Gonadotropin β subunits determine the rate of assembly and the oligosaccharide processing of hormone dimer in transfected cells. *J Cell Biol* 104:1173–1181
55. Currie RJ, McNeilly AS 1995 Mobilization of LH secretory granules in gonadotrophs in relation to gene expression, synthesis and secretion of LH during the preovulatory phase of the sheep oestrous cycle. *J Endocrinol* 147:259–270
56. Crawford JL, McNeilly JR, Nicol L, McNeilly AS 2002 Promotion of intragranular co-aggregation with LH by enhancement of secretogranin II storage resulted in increased intracellular granule storage in gonadotrophs of GnRH-deprived male mice. *Reproduction* 124:267–277
57. Crawford JL, McNeilly AS 2002 Co-localisation of gonadotropins and granins in gonadotrophs at different stages of the oestrous cycle in sheep. *J Endocrinol* 174:179–194
58. Crawford JL, McNeilly JR, McNeilly AS 2004 No evidence for pituitary priming to gonadotropin-releasing hormone in relation to luteinizing hormone (LH) secretion prior to the preovulatory LH surge in ewes. *Biol Reprod* 71:224–235
59. Thomas SG, Clarke IJ 1997 The positive feedback action of estrogen mobilizes LH-containing, but not FSH-containing secretory granules in ovine gonadotrophs. *Endocrinology* 138:1347–1350
60. Bhati M, Lee C, Nancarrow AL, Lee M, Craig VJ, Bach I, Guss JM, Mackay JP, Matthews JM 2008 Implementing the LIM code: the structural basis for cell type-specific assembly of LIM-homeodomain complexes. *EMBO J* 27:2018–2029
61. Eeckhoutte J, Briche I, Kurowska M, Formstecher P, Laine B 2006 Hepatocyte nuclear factor 4 α ligand binding and F domains mediate interaction and transcriptional synergy with the pancreatic islet LIM HD transcription factor Isl-1. *J Mol Biol* 364:567–581
62. Peng SY, Wang WP, Meng J, Li T, Zhang H, Li YM, Chen P, Ma KT, Zhou CY 2005 ISL-1 physically interacts with BETA2 to promote insulin gene transcriptional synergy in non- β cells. *Biochim Biophys Acta* 1731:154–159
63. West BE, Parker GE, Savage JJ, Kiratipranon P, Toomey KS, Beach LR, Colvin SC, Sloop KW, Rhodes SJ 2004 Regulation of the follicle-stimulating hormone β gene by the LHX3 LIM-homeodomain transcription factor. *Endocrinology* 145:4866–4879
64. Vasilyev VV, Lawson MA, Dipaolo D, Webster NJ, Mellon PL 2002 Different signaling pathways control acute induction versus long-term repression of LH β transcription by GnRH. *Endocrinology* 143:3414–3426
65. Tremblay JJ, Drouin J 1999 Egr-1 is a downstream effector of GnRH and synergizes by direct interaction with Ptx1 and SF-1 to enhance luteinizing hormone β gene transcription. *Mol Cell Biol* 19:2567–2576
66. Lamba P, Santos MM, Philips DP, Bernard DJ 2006 Acute regulation of murine follicle-stimulating hormone β subunit transcription by activin A. *J Mol Endocrinol* 36:201–220
67. Harrison GS, Wierman ME, Nett TM, Glode LM 2004 Gonadotropin-releasing hormone and its receptor in normal and malignant cells. *Endocr Relat Cancer* 11:725–748
68. Reis FM, Luisi S, Carneiro MM, Cobellis L, Federico M, Camargos AF, Petraglia F 2004 Activin, inhibin and the human breast. *Mol Cell Endocrinol* 225:77–82
69. Carro E, Pinilla L, Seoane LM, Considine RV, Aguilar E, Casanueva FF, Dieguez C 1997 Influence of endogenous leptin tone on the estrous cycle and luteinizing hormone pulsatility in female rats. *Neuroendocrinology* 66:375–377
70. Tena-Sempere M, Pinilla L, González LC, Diéguez C, Casanueva FF, Aguilar E 1999 Leptin inhibits testosterone secretion from adult rat testis *in vitro*. *J Endocrinol* 161:211–218
71. Lujan ME, Krzemien AA, Reid RL, Van Vugt DA 2006 Effect of leptin administration on ovulation in food-restricted rhesus monkeys. *Neuroendocrinology* 84:103–114
72. Di Yorio MP, Bilbao MG, Pustovrh MC, Prestifilippo JP, Faletti AG 2008 Leptin modulates the expression of its receptors in the hypothalamic-pituitary-ovarian axis in a differential way. *J Endocrinol* 198:355–366
73. Parent AS, Lebrethon MC, Gérard A, Vandersmissen E, Bourguignon JP 2000 Leptin effects on pulsatile gonadotropin releasing hormone secretion from the adult rat hypothalamus and interaction with cocaine and amphetamine regulated transcript peptide and neuropeptide Y. *Regul Pept* 92:17–24
74. Peyon P, Vega-Rubín de Celis S, Gómez-Requeni P, Zanuy S, Pérez-Sánchez J, Carrillo M 2003 *In vitro* effect of leptin on somatolactin release in the European sea bass (*Dicentrarchus labrax*): dependence on the reproductive status and interaction with NPY and GnRH. *Gen Comp Endocrinol* 132:284–292
75. Walczewska A, Yu WH, Karanth S, McCann SM 1999 Estrogen and leptin have differential effects on FSH and LH release in female rats. *Proc Soc Exp Biol Med* 222:170–177
76. Guo Z, Jiang H, Xu X, Duan W, Mattson MP 2008 Leptin-mediated cell survival signaling in hippocampal neurons mediated by JAK STAT3 and mitochondrial stabilization. *J Biol Chem* 283:1754–1763
77. Tartaglia LA 1997 The leptin receptor. *J Biol Chem* 272:6093–6096
78. Bates SH, Stearns WH, Dundon TA, Schubert M, Tso AW, Wang Y, Banks AS, Lavery HJ, Haq AK, Maratos-Flier E, Neel BG, Schwartz MW, Myers Jr MG 2003 STAT3 signalling is required for leptin regulation of energy balance but not reproduction. *Nature* 421:856–859
79. Hao A, Novotny-Diermayr V, Bian W, Lin B, Lim CP, Jing N, Cao X 2005 The LIM/homeodomain protein Islet1 recruits Janus tyrosine kinases and signal transducer and activator of transcription 3 and stimulates their activities. *Mol Biol Cell* 16:1569–1583
80. Takahashi Y, Okimura Y, Mizuno I, Iida K, Takahashi T, Kaji H, Abe H, Chihara K 1997 Leptin induces mitogen-activated protein kinase-dependent proliferation of C3H10T1/2 cells. *J Biol Chem* 272:12897–12900
81. Smith JT, Waddell BJ 2003 Developmental changes in plasma leptin and hypothalamic leptin receptor expression in the rat: peripubertal changes and the emergence of sex differences. *J Endocrinol* 176:313–319
82. Ketelslegers JM, Hetzel WD, Sherins RJ, Catt KJ 1978 Developmental changes in testicular gonadotropin receptors: plasma gonadotropins and plasma testosterone in the rat. *Endocrinology* 103:212–222